



DELAWARE HEALTH AND SOCIAL SERVICES

Division of Public Health

**Guidelines for the Management of
Multidrug Resistant and Other
Epidemiologically Important Organisms
along the Health Care Continuum**

**Delaware Health and Social Services
Division of Public Health
January 2010**

Guideline for the Management of Multidrug Resistant and Other Epidemiologically Important Organisms along the Health Care Continuum

Table of Contents

Topic	Page
Hospital Acquired Infections Advisory Sub-Committee Members	3
Hospital Acquired Infections Advisory Committee Members	4
Introduction	5
DPH Position Statement	5
Background	6
Risk Factors for MDROs	7
Diagnostic Testing	7
Infection Control Measures	7
Standard Precautions	8
Hand Hygiene	8
Respiratory Hygiene/Cough Etiquette	9
Room Placement	9
Group Activities	10
Environmental Cleaning	10
Shared Bathrooms, Showers, Tubs	11
Dishes, Glasses, Cups and Eating Utensils	11
Laundry	11
Decolonization	11
Indications for Contact Precautions	11
Contact Precautions: Gloves	12
Contact Precautions: Gowns	12
Contact Precautions: Patient Care Equipment	13
Masks, Eye Protection, Face shields	13
Discontinuing Contact Precautions	13
Communication with Other Facilities	13
Staff Education	13
Resident, Family, Visitor Education	14
Surveillance	14
Discharge Home	14
Antimicrobial Use	15
Summary	16
Abbreviations	16
Definitions	16
Attachment 1 (Standard Precautions)	19
References	21

Available on the DPH website at: <http://www.dhss.delaware.gov/dph/>

Hospital Acquired Infections Advisory Subcommittee Members

Donna Anderson RN.CIC.
Infection Preventionist
Beebe Medical Center

Michele Dennis, RN, BSN, MJ
VP of Nursing, Delaware Health Care Facilities Association, Board of Directors
Director of Nursing, Stonegates Health Center

Marci Drees, MD, MS, FACP
Hospital Epidemiologist & Medical Director, Infection Control
Christiana Care Health System

Brenda Johnson MT.CIC
Infection Control Practitioner
Nanticoke Memorial Hospital

Lawrence Markman, MD
DHSS

Jeanmarie McKinney, MSN, BSN, RN, CLNC
Quality Assurance Administrator
Division of Long Term Care Residents Protection
Delaware Health and Social Services

Yrene E. Waldron, LHNA
Executive Director
Delaware Health Care Facilities Association

Kathy Wroten, RN, BSN, CIC
Manager, Infection Control
Christiana Care Health System

Healthcare Associated Infections Advisory Committee Members

Kathy Wroten, RN, BSN, CIC (Chair)
Manager, Infection Control
Christiana Care Health System

Carol Briody MT, ASCP. CIC
(Co-Chair)
Infection Control Practitioner
Christiana Care Health System

Donna Anderson RN. CIC.
Infection Preventionist
Beebe Medical Center

Amanda Bundek MT.
Infection Control Practitioner
St. Francis Hospital

Marci Drees, MD, MS, FACP
Hospital Epidemiologist & Medical Director,
Infection Control
Christiana Care Health System

Kelly Gardner RN. CIC
Manager, Infection Control
Bayhealth Hospitals

Joanne Hasse
Public Representative

Brenda Johnson MT. CIC
Infection Control Practitioner
Nanticoke Memorial Hospital

Joel Klein, MD
Infectious Diseases Physician
Nemours/A.I. DuPont Hospital for Children

Sara Myers RN. BSN.
Nurse Trainer/Educator
Bureau of Correctional
Healthcare Services

Paul Silverman, Dr PH.
Associate Deputy Director
Delaware Division of Public Health

Jean Stipe RN
Infection Control Practitioner
Wilmington VA Medical Ctr.

Ramesh Vemulapalli, MD
Infectious Diseases Physician
Private Practice

Yrene E. Waldron, LHNA
Executive Director
Delaware Health Care
Facilities Association

Robert White
Health Insurer

Acknowledgement:

The Advisory Group would like to thank and acknowledge the authors of the "Guideline for the Management of Antimicrobial Resistant Microorganisms in Minnesota Long-Term Care Facilities,"¹ produced by the Minnesota Department of Health in October 2000 (revised 2004). This document was revised, updated and expanded by the Advisory Group to develop the Delaware guidelines.

Introduction:

The Delaware Division of Public Health (DPH) receives many inquiries from non-acute, residential care facilities along the health care continuum, including but not limited to nursing homes, rehabilitation centers, psychiatric and assisted living facilities, regarding the appropriateness of placement and care of persons with multidrug resistant organisms (MDRO). The most common questions concern persons who are colonized or infected with methicillin resistant *Staphylococcus aureus* (MRSA) or vancomycin resistant enterococcus (VRE), and increasingly those with highly resistant Gram-negative bacilli (GNB), such as *Pseudomonas*, *Klebsiella*, and *Acinetobacter* species. Due to concern about MDRO, some facilities have restricted the admission of persons known to be infected or colonized with MDRO, or have required negative screening cultures prior to transfer.² This has caused persons with MDRO to experience delays or denial of admission to long term care, psychiatric and other facilities.³ There is no evidence that restricting the admission of persons with MDRO is effective in keeping such facilities free of MDRO. Because residents are not generally screened for MDRO, colonization is often not detected. Therefore, a policy restricting the admission of persons with MDRO may lead to a false sense of security that a facility is free of MDRO.⁴ In addition, there is often concern regarding the transfer of patients recovering from transmissible gastrointestinal infections, particularly *Clostridium difficile* and Norovirus (Norwalk-like virus).

This guideline is intended to address all of the above “epidemiologically important” organisms, which for the purposes of this document will be collectively referred to using the general term MDRO. Guidance that is pertinent only to specific organisms will be designated as such. In addition, the term extended care facilities (ECF) will refer to all non-acute, residential care facilities along the health care continuum, including but not limited to nursing homes, rehabilitation centers, psychiatric and assisted living facilities.

DPH Position Statement:

Based on currently available knowledge from the Centers for Disease Control and Prevention (CDC), the Society for Healthcare Epidemiology of America (SHEA), the Association of Professionals in Infection Control and Hospital Epidemiology (APIC), and other sources, persons with MDRO should not be denied admission solely on the basis of a positive MDRO culture from any site, past history of any MDRO, or because they have recently been diagnosed with an MDRO.⁴⁻⁶ Also, it is not appropriate for ECFs to refuse to re-admit residents who have been found to have MDRO after transfer from the ECF to an acute care facility. New or returning residents should be admitted to ECFs based on the ability of the facility to provide necessary care to the resident and should not be based on MDRO status. Denying admission or re-admission based on MDRO status alone is discriminatory, may lead to review by government agencies, and is in violation of 16 Del. C., 1121(18). The CDC, SHEA, APIC, the American Hospital Association (AHA) task force and the Veterans’ Affairs (VA) consensus panel all oppose restricting the access of MRSA colonized residents to ECF.⁶⁻⁸ Guidelines published by other states also oppose such restrictions.⁹⁻¹¹

ECFs should be prepared to implement appropriate infection control measures for all prospective or current residents colonized or infected with MDRO. This guideline, which is based on the best current recommendations, describes infection control measures for ECF residents with MDRO in non-outbreak situations. It is important to note that due to

incomplete data specific to certain issues in the community extended care setting, extrapolation from studies done in other settings and/or situations has been necessary. As additional research is published, this guideline will be revised as necessary.

Background:

- The presence of MDRO colonization in ECF residents appears to be quite common. Studies have estimated that more than 20% of ECF residents may be colonized with MRSA and more than 10% may be colonized with VRE.¹²⁻¹⁸ In addition to MRSA and VRE, antibiotic resistant gram-negative bacilli (GNB) are also common in this setting (e.g., *E. coli*, *Pseudomonas aeruginosa*, *Klebsiella* species, and *Acinetobacter* species).^{15, 16, 19} In addition, increasing numbers of *C. difficile* infections have also been reported.²⁰ It has been shown that a significant portion of residents are already colonized with MRSA and VRE upon admission to ECFs and fewer residents acquire MRSA or VRE during their ECF stay.^{12, 21} The admission of persons with MDRO into ECFs does not appear to increase ECF infection rates or commonly result in excess morbidity or mortality.²²
- Although MDRO colonization may result in infection, it has been demonstrated that ECF residents colonized with MRSA and VRE are less likely to develop infections due to these organisms than are hospitalized patients.²³ Despite more frequent colonization, infections were less commonly due to MRSA or VRE than resistant GNB.²⁴ In studies of VA-affiliated ECF, few deaths resulted from MRSA infection and in a study of VRE colonization in an ECF, subsequent development of infection was rare.¹²⁻¹⁴ One study indicated that patients colonized with MRSA had higher 3-year mortality than non-colonized patients, but only among those with severe cognitive impairment.²⁵ A recent point-prevalence study of VA nursing homes demonstrated a 5.2% prevalence of nursing home-associated infections, with residents with any indwelling device having a 3-fold greater risk of infection.²⁶ Studies involving MRSA and VRE indicate that roommate to roommate transmission of these MDROs in the ECF setting, although possible, is not common.^{14, 27-29} Therefore, control measures that may be appropriate in the acute care setting are not generally indicated in the ECF setting. As in acute care hospitals, however, person-to-person spread via direct contact, particularly between residents and the transiently colonized hands of health care workers, is considered the principal mode of transmission.⁴ The importance of hand hygiene cannot be underemphasized in either the acute care or ECF setting.
- The prevalence of *C. difficile* in ECF in the absence of a recognized outbreak has ranged from 4 – 20%. Between 10 – 20% may acquire *C. difficile* during their stay. However, most residents may remain asymptomatic.³⁰ Viral gastroenteritis (caused by rotavirus, enteroviruses or norovirus) and bacterial gastroenteritis (including *C. difficile*) are well-known causes of diarrhea outbreaks in ECF. The elderly in particular are at increased risk for infectious gastroenteritis because of age-related decreases in gastric acid. Person-to-person spread, especially when bathrooms, dining and rehabilitation facilities are shared, plays an important role in viral gastroenteritis and *C. difficile* transmission.³¹ However, while patients recovering from these infections may continue to shed organisms in their feces,³² the cessation of diarrhea significantly decreases their infectivity to others.³¹

Risk Factors for MDROs:

The following factors have been found to be associated with MDRO colonization: underlying illness; intravenous, urinary, or enteral feeding devices; antibiotic use; wounds; previous hospitalizations; decline in functional status; advanced dementia; and increased intensity of nursing care.¹⁶⁻¹⁹ Factors that favor the spread of MDROs include high resident to staff ratios, lack of attention to basic infection control measures, use of common equipment without disinfection between residents, limited facilities for handwashing and the inappropriate use of antimicrobials.^{6, 23} ECFs must follow acceptable infection control standards of practice as indicated by the CDC, APIC, SHEA, and state health authorities.

Diagnostic Testing:

While active surveillance cultures have been recommended for acute care hospitals to limit spread of MRSA and/or VRE,³³⁻³⁶ this practice remains controversial^{33, 37-39} and active surveillance in ECF is not recommended.²⁸ Requiring negative screening cultures for MDRO prior to acceptance of patients from acute care settings leads to a false sense of security for several reasons:

- (1) Imperfect sensitivity of screening cultures, particularly if only one site (i.e., the nares for MRSA) is sampled;¹⁸
- (2) Increasing prevalence of community-associated MRSA in patients admitted directly from the community or from acute-care settings, who may never have had this colonization detected;^{23, 40}
- (3) Colonization may be intermittent,¹⁷ resulting in a patient whose initial screening cultures were negative but later becomes positive (but is undetected), particularly if the patient is placed on antimicrobial therapy.

It is not appropriate to require or request follow-up cultures of patients who have recovered from *C. difficile* or norovirus infection (i.e., a “test-of-cure”) and stool samples should never be sent for these pathogens from asymptomatic patients.³¹ If a patient with a history of *C. difficile* develops diarrhea after transfer to the ECF, it would be appropriate at this time to send an unformed stool sample for analysis, as recurrent disease is common. An unformed stool is defined as one which takes the shape of the container it is placed in. Formed stool samples should never be sent for *C. difficile* testing.

Infection Control Measures:

In 2006 the CDC published the guideline “Management of Multidrug-Resistant Organisms in Healthcare Settings.”²³ This guideline expanded on previous similar guidelines in part by incorporating recommendations not only for acute care facilities but also including ECF. This guidance acknowledged that “all healthcare settings are affected by the emergence and transmission of antimicrobial-resistant microbes” but that institutions “vary widely in physical and functional characteristics....Because of this, the approaches to prevention and control of these pathogens need to be tailored to the specific needs of each population and individual institution.” Modification of the recommendations is encouraged if the principles of epidemiology and disease transmission are maintained, and precautions are included to interrupt spread of infection by all routes that are likely to be encountered. In addition, the CDC published the “Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings 2007,”²⁹ which also incorporated recommendations for non-acute care settings. Information from these guidelines is summarized below.

Standard Precautions:

- Standard Precautions are based on the principle that all blood, body fluids, secretions, excretions (except sweat), non-intact skin, and mucous membranes may contain transmissible infectious agents. Standard Precautions include a group of infection prevention practices that apply to all patients, regardless of suspected or confirmed infection status, in any setting in which healthcare is delivered. These include: hand hygiene; use of gloves, gown, mask, eye protection, or face shield, depending on the anticipated exposure; and respiratory hygiene/cough etiquette.²⁹
- Most ECF residents with MDROs can be cared for using Standard Precautions (Attachment 1). Unless otherwise indicated (see below), Standard Precautions are adequate for ECF residents who have contained MDRO colonized or infected secretions/excretions.
 - Secretions/excretions (including wound drainage, stool and urine) are contained when they are unable to leak out of containment products. This includes wound dressings, incontinence products, urine bags, ostomy bags, etc.
- Because colonization with MDROs is often unrecognized and because other infectious organisms may also spread in ECFs, it cannot be emphasized too strongly that adherence with Standard Precautions is important in the care of all residents, whether or not they are known to have an MDRO. Emphasis should be placed on appropriate glove use and hand hygiene in ECF.^{6, 23, 29}

Hand Hygiene:

- MDROs are transmitted primarily via the contaminated hands of staff. The *single most effective means* of reducing the potential for MDRO transmission is hand hygiene (destroying or removing transient microorganisms from the hands) before and after contact with residents with MDROs, including after glove removal.
- Hand hygiene can be accomplished by cleaning hands with antimicrobial soap or by using a waterless alcohol-based hand antiseptic.^{41, 42} The use of a waterless alcohol-based hand antiseptic should be *strongly considered* in the care of all ECF residents. Recent studies show that such products are as effective as antimicrobial soaps, may increase handwashing compliance, and are not harmful to hands.⁴³⁻⁴⁵
- If hands are not visibly soiled and a waterless alcohol based product is to be used for hand antisepsis, apply a sufficient amount of the product to wet hands thoroughly, rub hands together and cover entire surface of hands (including nails) and allow to dry. However, if hands are visibly soiled with organic material or debris (e.g., feces, dirt), hands should be cleaned with soap and running water, rather than or prior to using a waterless product.
- Convenient access to hand hygiene products will improve adherence, therefore it is helpful to install pump dispensers of waterless alcohol-based hand antiseptic in or near each resident room. Small containers of waterless alcohol-based hand antiseptic may also be carried by staff and residents for use when handwashing facilities or pump dispensers are not readily accessible. This may be particularly useful for facilities/areas where placement of pump dispensers of alcohol-based products is not appropriate, such as psychiatric and dementia units.
- Caregivers should be cognizant of providing hand hygiene to the dependent resident population, especially before meals and after toileting, and should ensure educational efforts are provided for the residents in this area.

- For patients with *C. difficile* infection, alcohol-based products are not effective in removal of spores, although they will destroy any vegetative organisms present. Preferential use of soap and water instead of alcohol-based products for handwashing is a recommended consideration for outbreak situations.³¹ Glove use has been shown to significantly reduce the spread of *C. difficile* in hospitals.⁴⁶

Respiratory Hygiene/Cough Etiquette:²⁹

- This strategy is targeted at patients and accompanying family members and friends with undiagnosed transmissible respiratory infections, and applies to any person with signs of illness including cough, congestion, rhinorrhea, or increased production of respiratory secretions when entering a healthcare facility
- The elements of Respiratory Hygiene/Cough Etiquette include 1) education of healthcare facility staff, patients, and visitors; 2) posted signs, in language(s) appropriate to the population served, with instructions to patients and accompanying family members or friends; 3) source control measures (e.g., covering the mouth/nose with a tissue when coughing and prompt disposal of used tissues, using surgical masks on the coughing person when tolerated and appropriate); 4) hand hygiene after contact with respiratory secretions; and 5) spatial separation, ideally >3 feet, of persons with respiratory infections in common waiting areas when possible.
- Healthcare personnel are advised to observe Droplet Precautions (i.e., wear a mask) and perform hand hygiene when examining and caring for patients with signs and symptoms of a respiratory infection. Healthcare personnel who have a respiratory infection are advised to avoid direct patient contact, especially with high risk patients. If this is not possible, a mask should be worn while providing patient care.

Room Placement:

Residents with MDROs are not required to be placed in a private or single room. Single rooms may be limited in number and should be prioritized for those who have conditions that pose a risk of transmission to other residents/roommates either due to the specific MDRO or the ability to contain the infectious material (e.g., draining wounds, uncontained secretions). Single rooms are preferred for residents requiring Contact (see pg. 11) or Droplet precautions²⁹ and for patients with active *C. difficile* infection and fecal incontinence. If available, a single room should be considered for all *C. difficile* patients until diarrhea resolves.³⁰ Patients transferred from acute care facilities after treatment for an episode of *C. difficile* infection do not require private rooms or cohorting, presuming diarrhea is resolved.

If a single room is not available, residents with MDROs may be placed with appropriate roommates.²⁹ An appropriate roommate is either a resident with the same MDRO (cohorting) **or** a resident who:

- has intact skin with no significant open wounds or breaks in skin (superficial tears or breaks in the skin such as minor scratches, would be acceptable) *and*
- has no invasive devices, indwelling vascular or urinary catheters or drainage devices (e.g., tracheostomy or tracheal tubes, chest tubes, gastrostomy tubes, Broviac/Hickman catheters, intravenous lines, etc.) *and*
- is not significantly immunocompromised (e.g., due to organ transplantation, neutropenia, serious acute or chronic infection, systemic steroids or chemotherapy)
 - please note that based on age alone, most “healthy” elderly residents are not considered significantly immunocompromised – and is not colonized or infected with a different MDRO.

Contact DPH's Bureau of Epidemiology at 1-888-295-5156 if assistance is needed with placement issues. Situations in which a current resident is found to have an MDRO and there is concern about risk in a roommate should be evaluated on a case by case basis.

Group Activities:

Residents with MDROs may use common living areas, recreational areas, and group dining facilities. Such activities serve an important purpose in maintaining quality of life for ECF residents. Control measures that limit resident activity and movement, such as those used in the acute care setting, are generally not necessary and may result in emotional and social deprivation in the ECF setting.^{4, 6} The following factors should be considered in terms of group activities:

- Hand antisepsis is a very important component of participation in group activities. Residents with MDROs should have their hands cleaned with soap and water or a waterless alcohol based hand antiseptic prior to leaving their room and whenever they again become contaminated when the resident is out of their room.
- Residents with MDROs should have clean, dry wound dressings that adequately contain any drainage. Residents with MDROs in stool or urine, who are incontinent of either, should be clean and wear an incontinence product when leaving their room. In addition, all residents with MDROs should wear clean clothes or a clean cover gown when leaving their rooms.
- Residents with MDROs who are cognitively or behaviorally impaired and cannot maintain hygienic practices present challenges for ECFs. However, in most cases strategies can be devised that will address infection control issues as well as allow the resident the opportunity for movement and socialization. In rare cases some restriction of movement may be necessary (e.g., residents who may be shedding large numbers of MDRO bacteria and who have been linked to the development of infection in other residents). To discuss specific situations, please contact the DPH Bureau of Epidemiology at 1-888-295-5156.

Environmental Cleaning:

While the immediate patient environment (e.g., bed rails and tables) may become contaminated with MDRO, environmental cultures from common patient areas, including dining rooms, therapy areas, etc., are seldom positive.²⁸ Room cleaning - Standard facility procedures can be followed for cleaning the rooms of residents with MDRO in the ECF setting. Use of the facility's standard disinfectant is adequate, but it is important to monitor compliance with manufacturer's recommended application procedures and contact times. *C. difficile* spores and norovirus are relatively resistant to standard disinfection. Facilities may consider use of sodium hypochlorite (bleach)-containing cleaning agents for rooms housing patients with these organisms, particularly if the facility is experiencing high rates of infection that have not been controlled using standard means.^{30, 31}

- **Physical and recreational therapy equipment** - The hands of residents with MDROs should be cleaned before the resident uses recreational or physical therapy equipment. Standard facility procedures should be followed for routine cleaning and disinfection of recreational and physical therapy equipment used by residents with MDROs. However, this equipment should be disinfected before use by another resident if it becomes soiled with body secretions/excretions.
- **Trash disposal** - Follow standard facility procedures for trash and medical waste disposal. No special handling is necessary.²⁹

Shared Bathrooms, Showers, Tubs:

As previously stated, roommate to roommate transmission does not appear to be common in ECFs. Hygienic toileting practices (including thorough hand cleaning) are important for all residents, especially those dependent upon staff.

- **Bathrooms** - In situations where a resident with an MDRO shares a bathroom with a roommate who does not have the same MDRO, the bathroom should be cleaned and disinfected using standard facility procedure (e.g., daily and when visibly soiled). Commodes may be useful for certain residents with MDROs and should not be shared with roommates who do not have MDROs.
- **Showers, Tubs** - Shared tubs and showers should be cleaned and disinfected per standard facility procedure after use by residents with MDROs. It may be practical to bathe residents with MDROs after other residents.

Dishes, Glasses, Cups and Eating Utensils:

No special precautions are needed for dishes, glasses, cups or eating utensils. The combination of hot water and detergents used in institutional dishwashers is sufficient to decontaminate these items. All residents should be discouraged from sharing any of these items.

Laundry:

Standard precautions are adequate for handling laundry from all residents. However, soiled laundry, especially bed linens and towels from residents with uncontained stool, urine or other secretions/excretions, should be handled in such a way as to minimize contamination of staff and the environment. Gloves and long sleeved gowns should be worn when changing the beds of residents with uncontained stool, urine or other secretions/excretions and all soiled laundry should be placed directly in a moisture resistant container and not on the floor or other room surfaces. Special handling (i.e., double bagging, etc.) is not necessary. Laundry should not be rinsed at point of use. Laundry staff should wear gloves and long sleeved gowns when sorting laundry. No special laundering procedure is required.

Decolonization:

Routine decolonization for MRSA is not recommended for ECF residents.^{4, 23} Decolonization should not be required prior to admittance to an ECF.⁴ Decolonization therapy for MRSA may result in the emergence of resistance to the agents used and because recolonization is common, decolonization has had little impact on the incidence of infections experienced by ECF residents. There is no proven decolonization regimen for VRE, resistant GNB, or *C. difficile*. Among ECF residents, VRE colonization is likely to persist for extended periods of time and perhaps indefinitely.¹²

Indications for Contact Precautions (in addition to Standard Precautions):

Although Standard Precautions and hand antisepsis are sufficient for most residents with MDROs, Contact Precautions (see below) may be indicated for residents with MDROs who are potentially more likely to shed MDRO bacteria into their environment. A private room for ECF residents on Contact Precautions is recommended if available,⁶ but is not required.

The components of Contact Precautions that should be emphasized in this population are glove use and hand antisepsis. Residents for whom Contact Precautions, in addition to Standard Precautions, are indicated include the following:

- Residents who have MDRO infected or colonized wounds that cannot be covered fully by dressings or who have drainage that cannot be contained by dressings.

- Residents with fecal or urinary carriage of MDROs whose urine or stool cannot be contained in incontinence products, urine bags or ostomy bags.
- Residents with a tracheostomy who have an MDRO colonized or infected respiratory tract and large amounts of uncontained respiratory secretions.
- Residents who are ill and totally dependent on health care workers for their activities of daily living.
- Residents who have been epidemiologically linked to MDRO infections in other residents (or additional residents if an MDRO outbreak occurs).

Contact Precautions: Gloves

- In addition to wearing gloves as necessary for Standard Precautions, gloves (clean, nonsterile gloves are adequate) should be worn when providing direct care (changing clothing, toileting, bathing, dressing changes, etc.) to residents with MDROs.
- Gloves should also be worn when handling items potentially contaminated by MDROs. Potentially contaminated items include bedside tables, overbed tables, bed rails, bathroom fixtures, television and bed controls, IV poles, suction and oxygen tubing and electronic equipment, especially control knobs. If urine or stool is colonized or infected with an MDRO, gloves should be worn when handling urine/stool containers or bags.
- During the course of providing care for residents, gloves should be changed and hands cleaned after contact with material that could contain high concentrations of microorganisms (e.g., stool, urine or wound drainage).
- Wearing gloves is not a substitute for hand antisepsis. Gloves should be removed and discarded and hands cleaned (see Hand Antisepsis above) after providing care to a resident with an MDRO and before leaving the resident's immediate environment or providing care to another resident. Avoid touching items in the resident's room with contaminated gloves and avoid touching potentially contaminated surfaces or items after glove removal and hand antisepsis.

Contact Precautions: Gowns

- For all residents, gowns should be worn as necessary for Standard Precautions. In addition, when caring for residents on Contact Precautions clean, non-sterile gowns should be worn if direct care (e.g., bathing, lifting) will be provided or when substantial contact with secretions/excretions (e.g., linen changes) is anticipated. When such contact is anticipated it is generally most practical to put on a gown before entering the room or before approaching the resident. Long-sleeved gowns typically provide the best protection.
- Gowns should also be worn when contact with environmental surfaces and items in the resident's room which are likely to be contaminated (those close to or used by the resident) is anticipated. This is particularly true if the resident is incontinent of urine or stool or has diarrhea, an ileostomy, a colostomy, or wound drainage not contained by a dressing.
- Gowns should be removed and discarded before leaving the resident's environment. After gown removal, ensure that clothing does not contact potentially contaminated environmental surfaces. Gowns should not be reused, even by the same staff member, making multiple trips into the same room.

Contact Precautions: Patient Care Equipment

- Dedicate patient care equipment to a single resident (e.g., commodes, thermometers, blood pressure cuffs, and stethoscopes). If equipment is to be shared it must be cleaned and disinfected before use by another resident. Use of standard facility disinfectants is adequate. Personal items of residents with MDROs should not be shared with other residents unless they have been cleaned and disinfected. (See Environmental Cleaning)

Masks, Eye Protection, Face Shields:

MRSA, VRE, resistant GNB, and *C. difficile* are not known to be transmitted via the airborne route and masks are not a routine component of Contact Precautions. However, masks and eye protection or face shields should be worn during the care of any resident if indicated by Standard Precautions. Masks and eye protection or face shields should be worn during resident care activities that are likely to generate splashes or sprays of blood, body fluids, secretions or excretions. Such activities include suctioning or working within 3 feet of a resident who is likely to expel droplet secretions (e.g., those with uncontained respiratory secretions, those with tracheitis and a tracheostomy tube in place, or those with suspected norovirus infection who are actively vomiting).

Discontinuing Contact Precautions:

When the condition of a resident with an MDRO changes (e.g., drainage is contained) and they therefore no longer meet the criteria for needing Contact Precautions, they may be cared for using Standard Precautions. However, if the resident continues to have secretions or excretions that are not contained, Contact Precautions should be continued.

It is appropriate to obtain cultures for clinical reasons, such as determining if infection is present or to guide antimicrobial treatment. However, non-clinically indicated cultures to determine if a resident remains colonized with an MDRO are not generally recommended. It is likely that ECF residents who are colonized with MDROs will continue to be colonized for significant lengths of time and repeated culturing will not be beneficial.

Communication with Other Facilities:

It is important for ECF and acute care facilities to work together in an attempt to control the spread of MDROs. Effective communication between LTC and acute care facilities is important in ensuring that the MDRO status of residents is known and that appropriate precautions are instituted or maintained in both types of facilities. Because of the acuity level, the frequent presence of invasive devices, and the vulnerability of many patients in the acute care setting, Contact Precautions are generally implemented for all patients known to have MDROs in the acute care setting. Because Contact Precautions may not be indicated for the same resident in the ECF, the rationale for the difference in precautions (as described above) should be explained to the resident and their family. When a resident is discharged from an acute care facility or transferred from an ECF, an adequate history should be provided to ensure that the receiving facility can institute the proper precautions.

Staff Education:

Continuing education programs for staff who have direct contact with residents or the items in the residents' environment is strongly encouraged. Staff who are responsible for making decisions regarding the care of residents should receive information about MDROs and should thoroughly review the information presented in this guideline. When educating staff

about MDRO it should be noted that healthy people are at very little risk of developing an infection with MDRO. In addition, there are no special precautions for pregnant staff who work with residents with MDRO.

Resident, Family, Visitor Education:

- Residents and their families and visitors should be educated about the MDRO and the precautions to be taken in the ECF, including hand antisepsis and methods to limit environmental contamination with stool, urine, and respiratory secretions.
- Family and visitors should clean their hands before entering and leaving the room of a resident with an MDRO. Family/visitors should wear gloves when handling the secretions/excretions of residents with MDROs or when providing direct care (e.g., bathing). Hands should be cleaned after glove removal.
- Regardless of whether additional infection control precautions (Contact Precautions) are in place for a particular resident, it is important to encourage all family/visitors to clean their hands before and after visiting any resident.
- In addition, families, visitors and other residents must have sufficient education to alleviate their concerns, ensure that precautions are maintained, and understand that residents with MDROs need not be avoided.¹⁹

Surveillance:

- Surveillance is an important component of all infection control programs, including those in ECF.^{6, 29} Surveillance should include the regular review of all microbiology culture and susceptibility data to detect MDRO and other epidemiologically important organisms.^{4, 6, 29}
- Although residents known to have MRSA and VRE are likely to represent only a portion of the residents with MDRO, screening residents for MDRO is **not** indicated as a routine infection control measure in ECF.²⁹
- A confidential line list of residents with MDROs should be maintained.
- Surveillance of cultures obtained for clinical reasons can establish baseline rates of MDRO infections for a facility. High rates of endemic infection (e.g., more than 1 case per 1,000 resident-care days) or an outbreak (e.g., > 3 infections in one week or twice the number of infections per month observed for 3 consecutive months) may be identified by surveillance.^{6, 29}
- Facilities can use surveillance data in educational programs to reinforce infection control practices and to prioritize infection control activities.
- Culture and susceptibility data can be used to guide the choice of empiric therapy. Empiric therapy is the antimicrobial treatment that is prescribed based on a presumed diagnosis before culture and susceptibility results are known. Monitoring culture and susceptibility data will provide information on antimicrobial susceptibility and resistance patterns in a particular facility. Communicating this information can help guide physicians in the selection of the most appropriate empiric therapy for residents with infections.

Discharge Home:

- Family members should be advised that healthy persons have very little risk of developing an infection due to an MDRO.
- Family members should wear gloves when handling secretions/excretions that are known to contain MDROs and should clean their hands with soap and water or a waterless alcohol based hand antiseptic after glove removal. When gloves have not been used, family members should clean their hands after providing direct care.

- The resident's usual healthcare providers should be informed of the MDRO.
- The person with an MDRO should clean his/her hands frequently with soap and water or a waterless alcohol based hand antiseptic.
- No special laundering procedures are necessary. The clothes of all family members can be washed together using standard detergent. Bleach can be added for items that are heavily soiled with body secretions/excretions.
- Regular cleaning with household disinfectants of surfaces and items contaminated by secretions/excretions or frequently touched by hands is recommended.
- Family members of persons with MDRO should clean their hands with soap or a waterless alcohol based hand antiseptic before preparing food and before eating.
- Persons who are very ill or have compromised immune systems should avoid handling secretions/excretions known to contain MDRO. Such persons should also clean their hands with soap and water or a waterless alcohol based hand antiseptic after contact with a person with an MDRO.

Antimicrobial Use:

The appropriate use of antimicrobials is **the most important** method of controlling antimicrobial resistance. One of the ways resistance develops is through the use and overuse of antimicrobials (antimicrobial pressure). When bacteria are exposed to an antimicrobial, those bacteria that are susceptible to the antimicrobial are killed and those that are resistant survive and may become predominant. These antimicrobial resistant organisms may then be transmitted to other persons. Overuse of antimicrobials is a problem in ALL healthcare settings. Studies have repeatedly documented that much of the antimicrobial use in ECF is inappropriate.⁴⁷

Attempts to control antimicrobial use in ECF are complicated by many factors, including the fact that clinical diagnosis is often difficult in this population due to the frequent absence of typical signs and symptoms of infection (e.g., fever). However, ECFs should attempt to develop and institute programs and policies to monitor and control the use of antimicrobials in all residents. SHEA recently published a position paper on antimicrobial use in long-term

care facilities.⁴⁷ This position paper outlines concerns about inappropriate antimicrobial use and recommends approaches to promote the rational use of antimicrobials in this setting. The CDC has also published a guideline regarding the prudent use of vancomycin.⁴⁸

- In particular, treatment of asymptomatic bacteriuria is very common in ECF. Such treatment is often inappropriate and can induce the development of resistant organisms. More than 30% of non-catheterized residents and almost all chronically catheterized residents have asymptomatic bacteriuria.⁴⁷
- The use of antimicrobials for the treatment of viral upper respiratory infections is common and also considered to be inappropriate.⁴⁹
- Evaluation of the resident's clinical signs and symptoms in conjunction with appropriate culture and susceptibility testing should be used to guide therapy.⁴⁷
- Although empiric antimicrobial therapy may be indicated in some situations, it is important that physicians are informed of culture and susceptibility results so that applicable changes in treatment can be made. Broad spectrum antimicrobials (which are often used for empiric therapy) should be replaced by narrower spectrum antimicrobials once culture and susceptibility results are known. The use of narrower spectrum antimicrobials that target specific microorganisms will lessen antimicrobial pressure. In addition, stopping empiric antimicrobials promptly if no

bacterial infection is identified, and using antimicrobials for the shortest duration necessary to treat the infection (i.e., 7 – 8 days rather than 14 days for most healthcare-associated pneumonia),⁵⁰ are both important strategies to reduce antimicrobial pressure.

- Because residents with invasive devices are at greater risk of acquiring an MDRO, minimizing the use of invasive devices such as urinary catheters, feeding tubes, tracheostomies, etc. may be helpful in controlling MDROs in ECFs.⁴
- Avoiding antimicrobial use in patients with a history of *C. difficile* infection is particularly important, as repeated or continued antimicrobial use can lead to recurrent *C. difficile* infections that may be severe. For patients with a history of *C. difficile* who require antimicrobials, appropriate surveillance should be in place so that if diarrhea does recur, prompt diagnostic testing, infection control measures, and treatment if necessary can be instituted.

Summary:

This guideline was developed to assist Delaware ECF in achieving the goal of prevention and control of MDRO while maintaining quality of life for those residents who are colonized or infected with MDRO. Using this guideline as a reference, ECF will be able to assess individual residents with MDRO and develop specific care plans to minimize both the restrictions imposed on the resident and the risk of transmission of MDRO to others. It is important that the staff who develop care plans and those who implement them have sufficient knowledge to make appropriate decisions.

Please remember that the DPH Bureau of Epidemiology (1-888-295-5156) is available for consultation regarding questionable or difficult situations, including possible outbreaks, at any time. DPH does not report any issues related to MDRO to any state or other regulatory agency.

Abbreviations:

- **APIC** = Association of Professionals in Infection Control and Hospital Epidemiology
- **CDC** = Centers for Disease Control and Prevention
- **ECF** = extended care facilities
- **DPH** = Division of Public Health
- **GNB** = Gram-negative bacilli
- **MDRO** = multi-drug resistant organisms
- **MRSA** = methicillin resistant *Staphylococcus aureus*
- **SHEA** = Society for Healthcare Epidemiology of America
- **VRE** = vancomycin resistant Enterococcus

Definitions:

- ***Clostridium difficile* (C diff)**: anaerobic bacteria that are the most common cause of antibiotic-associated diarrhea. May lead to asymptomatic infection or cause varying severity of diarrhea, pseudomembranous colitis, toxic megacolon, and death.

- **Cohort:** two or more patients colonized or infected with a specific MDRO, who are physically separated from other patients who are not known to be colonized or infected with the same MDRO.
- **Colonization:** the presence of microorganisms in or on a host with growth and multiplication but without tissue invasion or damage (*vs. Infection*).
- **Contact Precautions (or Isolation):** Additional procedures put in place to reduce spread of microorganisms that are spread by direct contact with patients or their environment.
- **Decolonization:** the administration of topical, oral, or systemic antibiotics or antiseptics for the purpose of eradicating MDRO colonization.
- **Decontamination:** the process of removing microorganisms and making an object safe for handling.
- **Disinfection:** a process that kills or destroys nearly all microorganisms, with the exception of bacterial spores, on inanimate objects.
- **Droplet Precautions (or Isolation):** Additional procedures put in place to reduce spread of microorganisms that are spread by large respiratory particles produced by coughing or sneezing. These droplets typically do not persist in the air beyond 3 – 6 feet of the patient, and standard surgical masks are used to protect healthcare workers.
- **Enterococcus species** (includes *E. faecium* and *E. faecalis*): common gram-positive microorganisms that are part of the normal flora in the gastrointestinal and female genital tract.
- **Extended care facilities (ECF):** in this document, refers to all non-acute, residential care facilities along the health care continuum, including but not limited to nursing homes, rehabilitation centers, psychiatric and assisted living facilities.
- **Highly resistant Gram-negative bacteria (GNB):** a heterogenous group of Gram-negative bacteria that have acquired resistance to commonly used antimicrobials. Includes organisms that produce extended spectrum beta-lactamases (ESBL) and carbapenemases, which inactivate beta-lactam (such as penicillins and cephalosporins) and carbapenem antibiotics, respectively. Common organisms include but are not limited to *E. coli*, *Klebsiella*, *Acinetobacter*, and *Pseudomonas* species.
- **Infection:** the entry and multiplication of microorganisms in the tissues of the host accompanied by clinical signs and symptoms (*vs. Colonization*).
- **Methicillin Resistant *Staphylococcus aureus* (MRSA):** strains of *S. aureus* that are resistant to the antibiotics methicillin, oxacillin, nafcillin and frequently other antimicrobials.
- **Multidrug-resistant organisms (MDRO):** a term that is used collectively to refer to bacteria that have acquired resistance to 2 or more commonly used antimicrobials. Includes MRSA, VRE, and resistant GNB. For the purposes of this document, also refers to any epidemiologically important organism, such as *C. difficile* and norovirus, which are not technically multidrug-resistant.
- **Norovirus (also known as Norwalk-like viruses):** a type of calcivirus that causes highly contagious viral gastroenteritis characterized by nausea, vomiting and diarrhea that typically lasts 1-3 days.
- **Outbreak:** a definite increase in the number of new MDRO-infected patients above the endemic rate in a defined period of time (e.g., a month).
- **Prevalence:** the proportion of persons in the facility with MDRO during a specific time period.

- **Sanitation:** a process that decreases microorganisms on an inanimate object to a safe or relatively safe level.
- **Standard Precautions:** Procedures intended to be used for all patients to reduce spread of microorganisms (see Attachment 1).
- ***Staphylococcus aureus*:** a common species of gram-positive bacteria found on the skin and in the anterior nares of many people.
- **Surveillance:** monitoring the distribution of illnesses in a population.
- **Transmission:** the method or route by which microorganisms move from one location to another (e.g., by direct contact, airborne, droplet, or indirect contact with inanimate objects such as equipment).
- **Vancomycin Resistant Enterococcus (VRE):** strains of enterococcus (*E. faecalis* and *E. faecium*) that are resistant to the antibiotic vancomycin and other antimicrobial.

The Delaware Division of Public Health wishes to express appreciation to the members of the Advisory Group for their assistance in the development of this guideline.

For more information or additional copies, please call the DPH Bureau of Epidemiology at 1-888-295-5156. This document may also be downloaded at:
<http://www.dhss.delaware.gov/dph/>.

Attachment 1:

STANDARD PRECAUTIONS

From: **Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings 2007.**²⁹

RECOMMENDATIONS FOR APPLICATION OF STANDARD PRECAUTIONS FOR THE CARE OF ALL PATIENTS IN ALL HEALTHCARE SETTINGS

The application of Standard Precautions during patient care is determined by the nature of the health care worker-patient interaction and the extent of anticipated blood, body fluid, or pathogen exposure. For some interactions (e.g., performing venipuncture), only gloves may be needed; during other interactions (e.g., suctioning), use of gloves, gown, and face shield or mask and goggles is necessary.

COMPONENT RECOMMENDATIONS

A. Hand hygiene

- (1) After touching blood, body fluids, secretions, excretions, contaminated items; immediately after removing gloves; between patient contacts.
- (2) Wash hands regardless of whether gloves are worn.
- (3) It may be necessary to wash hands between tasks and procedures on the same patient to prevent cross-contamination of different body sites.

B. Personal protective equipment (PPE)

- (1) Gloves for touching blood, body fluids, secretions, excretions, contaminated items; for touching mucous membranes and non-intact skin
- (2) Gown during procedures and patient-care activities when contact of clothing/exposed skin with blood/body fluids, secretions, and excretions is anticipated..
- (3) Mask, eye protection (goggles), face shield* during procedures and patient-care activities likely to generate splashes or sprays of blood, body fluids, secretions, especially suctioning, endotracheal intubation

*During aerosol-generating procedures on patients with suspected or proven infections transmitted by respiratory aerosols (e.g., SARS), wear a fit-tested N95 or higher respirator in addition to gloves, gown, and face/eye protection.

C. Respiratory hygiene/cough etiquette (source containment of infectious respiratory secretions in symptomatic patients, beginning at initial point of encounter e.g., triage and reception areas in emergency departments and physician offices)

- (1) Instruct symptomatic persons to cover mouth/nose when sneezing/coughing
- (2) Use tissues and dispose in no-touch receptacle
- (3) Observe hand hygiene after soiling of hands with respiratory secretions
- (4) Wear surgical mask if tolerated or maintain spatial separation, >3 feet if possible.

D. Patient placement

- (1) Prioritize single-patient rooms for patients at increased risk of transmission, likely to contaminate the environment, does not maintain appropriate hygiene, or is at increased risk of acquiring infection or developing adverse outcome following infection.

E. Soiled patient-care equipment

(1) Handle in a manner that prevents transfer of microorganisms to others and to the environment; wear gloves if visibly contaminated; perform hand hygiene.

F. Environmental control

(1) Develop procedures for routine care, cleaning, and disinfection of environmental surfaces, especially frequently touched surfaces in patient-care areas.

G. Textiles and laundry

(1) Handle in a manner that prevents transfer of microorganisms to others and to the environment

H. Needles and other sharps

(1) Do not recap, bend, break, or hand-manipulate used needles; if recapping is required, use a one-handed scoop technique only

(2) Use safety features when available

(3) Place used sharps in puncture-resistant container

I. Patient resuscitation

(1) Use mouthpiece, resuscitation bag, other ventilation devices to prevent contact with mouth and oral secretions

References

1. Minnesota Department of Health, Infectious Disease Prevention and Control Division, Facility and Provider Compliance Division. Guideline for the management of antimicrobial resistant microorganisms in Minnesota long-term care facilities. October 2000. Available at:
<http://www.health.state.mn.us/divs/idepc/dtopics/infectioncontrol/ltcguideline.html>
2. Thurn JR, Crossley K, Gerdt A. Bacterial colonization of nursing home residents on admission to an acute care hospital. *J Hosp Infect.* 1996;32(2):127-133.
3. Bryce EA, Tiffin SM, Isaac-Renton JL, Wright CJ. Evidence of delays in transferring patients with methicillin-resistant staphylococcus aureus or vancomycin-resistant enterococcus to long-term-care facilities. *Infect Control Hosp Epidemiol.* 2000;21(4):270-271.
4. Strausbaugh LJ, Crossley KB, Nurse BA, Thrupp LD. Antimicrobial resistance in long-term-care facilities. *Infect Control Hosp Epidemiol.* 1996;17(2):129-140.
5. Smith PW, Bennett G, Bradley S, Drinka P, Lautenbach E, Marx J, Mody L, Nicolle L, Stevenson K. Society for Healthcare Epidemiology of America (SHEA). Association for Professionals in Infection Control and Epidemiology (APIC). SHEA/APIC guideline: Infection prevention and control in the long-term care facility. *Am J Infect Control.* 2008;36(7):504-535.
6. Smith PW, Bennett G, Bradley S, et al. SHEA/APIC guideline: Infection prevention and control in the long-term care facility, July 2008. *Infect Control Hosp Epidemiol.* 2008;29(9):785-814.
7. Boyce JM, Jackson MM, Pugliese G, et al. Methicillin-resistant staphylococcus aureus (MRSA): A briefing for acute care hospitals and nursing facilities. The AHA technical panel on infections within hospitals. *Infect Control Hosp Epidemiol.* 1994;15(2):105-115.
8. Mulligan ME, Murray-Leisure KA, Ribner BS, et al. Methicillin-resistant *Staphylococcus aureus*: A consensus review of the microbiology, pathogenesis, and epidemiology with implications for prevention and management. *Am J Med.* 1993;94(3):313-328.
9. Cahill CK, Rosenberg J. Guideline for prevention and control of antibiotic-resistant microorganisms in California long-term care facilities. Centers for Disease Control. *J Gerontol Nurs.* 1996;22(5):40-47.
10. Supplemental infection control guidelines for the care of patients colonized or infected with vancomycin-resistant enterococci (VRE) in hospitals, long-term care facilities and home health care, 2009. Available at:
http://www.health.state.ny.us/professionals/diseases/reporting/communicable/infection/95-14_vre_control_guidelines.htm. Accessed 10/8/2009.
11. Colorado Medical Directors Association, Colorado Department of Public Health and Environment. Recommendations for prevention and control of methicillin-resistant

Staphylococcus aureus in long-term-care facilities. June 2009. Available at: www.cdphe.state.co.us/hf/protocols/MRSA%20LTCFrecs%20final%206_05.pdf.

12. Bradley SF, Terpenning MS, Ramsey MA, et al. Methicillin-resistant *Staphylococcus aureus*: Colonization and infection in a long-term care facility. *Ann Intern Med*. 1991;115(6):417-422.
13. Strausbaugh LJ, Jacobson C, Sewell DL, Potter S, Ward TT. Methicillin-resistant *Staphylococcus aureus* in extended-care facilities: Experiences in a Veterans' Affairs nursing home and a review of the literature. *Infect Control Hosp Epidemiol*. 1991;12(1):36-45.
14. Bonilla HF, Zervos MA, Lyons MJ, et al. Colonization with vancomycin-resistant *Enterococcus faecium*: Comparison of a long-term-care unit with an acute-care hospital. *Infect Control Hosp Epidemiol*. 1997;18(5):333-339.
15. Furuno JP, Hebden JN, Standiford HC, et al. Prevalence of methicillin-resistant *Staphylococcus aureus* and *Acinetobacter baumannii* in a long-term acute care facility. *Am J Infect Control*. 2008;36(7):468-471.
16. Pop-Vicas A, Mitchell SL, Kandel R, Schreiber R, D'Agata EM. Multidrug-resistant Gram-negative bacteria in a long-term care facility: Prevalence and risk factors. *J Am Geriatr Soc*. 2008;56(7):1276-1280.
17. Stone ND, Lewis DR, Lowery HK, et al. Importance of bacterial burden among methicillin-resistant *Staphylococcus aureus* carriers in a long-term care facility. *Infect Control Hosp Epidemiol*. 2008;29(2):143-148.
18. Mody L, Kauffman CA, Donabedian S, Zervos M, Bradley SF. Epidemiology of *Staphylococcus aureus* colonization in nursing home residents. *Clin Infect Dis*. 2008;46(9):1368-1373.
19. Bradley SF. Issues in the management of resistant bacteria in long-term-care facilities. *Infect Control Hosp Epidemiol*. 1999;20(5):362-366.
20. Crnich CJ, Safdar N, Robinson J, Zimmerman D. Longitudinal trends in antibiotic resistance in US nursing homes, 2000-2004. *Infect Control Hosp Epidemiol*. 2007;28(8):1006-1008.
21. Brennen C, Wagener MM, Muder RR. Vancomycin-resistant *Enterococcus faecium* in a long-term care facility. *J Am Geriatr Soc*. 1998;46(2):157-160.
22. Spindel SJ, Strausbaugh LJ, Jacobson C. Infections caused by *Staphylococcus aureus* in a Veterans' Affairs nursing home care unit: A 5-year experience. *Infect Control Hosp Epidemiol*. 1995;16(4):217-223.
23. Siegel JD, Rhinehart E, Jackson M, Chiarello L, the Healthcare Infection Control Advisory Committee. Management of multi-drug resistant organisms in healthcare settings, 2006. *Am J Infect Control*. 2007 Dec;35(10 Suppl 2):S165-93.

24. Terpenning MS, Bradley SF, Wan JY, Chenoweth CE, Jorgensen KA, Kauffman CA. Colonization and infection with antibiotic-resistant bacteria in a long-term care facility. *J Am Geriatr Soc*. 1994;42(10):1062-1069.
25. Suetens C, Niclaes L, Jans B, et al. Methicillin-resistant *Staphylococcus aureus* colonization is associated with higher mortality in nursing home residents with impaired cognitive status. *J Am Geriatr Soc*. 2006;54(12):1854-1860.
26. Tsan L, Davis C, Langberg R, et al. Prevalence of nursing home-associated infections in the Department of Veterans Affairs nursing home care units. *Am J Infect Control*. 2008;36(3):173-179.
27. Hsu CC. Serial survey of methicillin-resistant *Staphylococcus aureus* nasal carriage among residents in a nursing home. *Infect Control Hosp Epidemiol*. 1991;12(7):416-421.
28. Nicolle LE. Infection control in long-term care facilities. *Clin Infect Dis*. 2000;31(3):752-756.
29. Siegel JD, Rhinehart E, Jackson M, Chiarello L, Health Care Infection Control Practices Advisory Committee. 2007 Guideline for isolation precautions: Preventing transmission of infectious agents in health care settings. *Am J Infect Control*. 2007;35(10 Suppl 2):S65-164.
30. Simor AE, Bradley SF, Strausbaugh LJ, Crossley K, Nicolle LE, SHEA Long-Term-Care Committee. *Clostridium difficile* in long-term-care facilities for the elderly. *Infect Control Hosp Epidemiol*. 2002;23(11):696-703.
31. Dubberke ER, Gerding DN, Classen D, et al. Strategies to prevent *Clostridium difficile* infections in acute care hospitals. *Infect Control Hosp Epidemiol*. 2008;29 Suppl 1:S81-92.
32. Siebenga JJ, Beersma MF, Vennema H, van Biezen P, Hartwig NJ, Koopmans M. High prevalence of prolonged norovirus shedding and illness among hospitalized patients: A model for in vivo molecular evolution. *J Infect Dis*. 2008;198(7):994-1001.
33. Farr BM. Political versus epidemiological correctness. *Infect Control Hosp Epidemiol*. 2007;28(5):589-593.
34. Coia JE, Duckworth GJ, Edwards DI, et al. Guidelines for the control and prevention of methicillin-resistant *Staphylococcus aureus* (MRSA) in healthcare facilities. *Journal of Hospital Infection*. 63 Suppl 1:S1-44, 2006 May. 2006:S1-44.
35. Farr BM. What to think if the results of the national institutes of health randomized trial of methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant enterococcus control measures are negative (and other advice to young epidemiologists): A review and an au revoir. *Infect Control Hosp Epidemiol*. 2006;27(10):1096-1106.
36. Muto CA, Jernigan JA, Ostrowsky BE, et al. SHEA guideline for preventing nosocomial transmission of multidrug-resistant strains of *Staphylococcus aureus* and enterococcus. *Infect Control Hosp Epidemiol*. 2003;24(5):362-386.

37. Muto CA. Methicillin-resistant *Staphylococcus aureus* control: We didn't start the fire, but it's time to put it out. *Infect Control Hosp Epidemiol*. 2006;27(2):111-115.
38. de San N, Denis O, Gasasira MF, De Mendonca R, Nonhoff C, Struelens MJ. Controlled evaluation of the IDI-MRSA assay for detection of colonization by methicillin-resistant *Staphylococcus aureus* in diverse mucocutaneous specimens. *J Clin Microbiol*. 2007;45(4):1098-1101.
39. Strausbaugh LJ, Siegel JD, Weinstein RA. Preventing transmission of multidrug-resistant bacteria in health care settings: A tale of 2 guidelines. *Clin Infect Dis*. 2006;42(6):828-835.
40. Weber JT. Community-associated methicillin-resistant *Staphylococcus aureus*. *Clin Infect Dis*. 2005;41 Suppl 4:S269-72.
41. Larson EL. APIC guideline for handwashing and hand antisepsis in health care settings. *Am J Infect Control*. 1995;23(4):251-269.
42. Boyce JM, Pittet D, Healthcare Infection Control Practices Advisory Committee, HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. Guideline for hand hygiene in health-care settings. recommendations of the healthcare infection control practices advisory committee and the HICPAC/SHEA/APIC/IDSA hand hygiene task force. Society for Healthcare Epidemiology of America/Association for Professionals in Infection Control/Infectious Diseases Society of America. *MMWR Recomm Rep*. 2002;51(RR-16):1-45.
43. Bischoff WE, Reynolds TM, Sessler CN, Edmond MB, Wenzel RP. Handwashing compliance by health care workers: The impact of introducing an accessible, alcohol-based hand antiseptic. *Arch Intern Med*. 2000;160(7):1017-1021.
44. Boyce JM. Using alcohol for hand antisepsis: Dispelling old myths. *Infect Control Hosp Epidemiol*. 2000;21(7):438-441.
45. Boyce JM, Kelliher S, Vallande N. Skin irritation and dryness associated with two hand-hygiene regimens: Soap-and-water hand washing versus hand antisepsis with an alcoholic hand gel. *Infect Control Hosp Epidemiol*. 2000;21(7):442-448.
46. Johnson S, Gerding DN, Olson MM, et al. Prospective, controlled study of vinyl glove use to interrupt *Clostridium difficile* nosocomial transmission. *Am J Med*. 1990;88(2):137-140.
47. Nicolle LE, Bentley DW, Garibaldi R, Neuhaus EG, Smith PW. Antimicrobial use in long-term-care facilities. SHEA long-term-care committee. *Infect Control Hosp Epidemiol*. 2000;21(8):537-545.

48. Recommendations for preventing the spread of vancomycin resistance: Recommendations of the hospital infection control practices advisory committee (HICPAC). *Am J Infect Control*. 1995;23(2):87-94.
49. Gonzales R, Steiner JF, Sande MA. Antibiotic prescribing for adults with colds, upper respiratory tract infections, and bronchitis by ambulatory care physicians. *JAMA*. 1997;278(11):901-904.
50. American Thoracic Society, Infectious Diseases Society of America. Guidelines for the management of adults with hospital-acquired, ventilator-associated, and healthcare-associated pneumonia. *Am J Respir Crit Care Med*. 2005;171(4):388-416.