

Delaware Weekly Influenza Report MMWR Week 15 (April 10 - 16, 2016) Delaware Division of Public Health

National Influenza Synopsis 2015-2016:

National data are updated Friday of each week. Please visit http://www.cdc.gov/flu/weekly/ for the most current information. During MMWR Week 15 (April 10 - 16, 2016) influenza activity decreased in the United States. The most frequently identified influenza virus type reported by public health laboratories during week 15 was influenza A, with influenza A (H1N1)pdm09 viruses predominating. The percentage of respiratory specimens testing positive for influenza in clinical laboratories decreased. Widespread influenza activity was reported by Puerto Rico and 14 states (Colorado, Connecticut, Delaware, Kentucky, Maine, Massachusetts, Nebraska, New Hampshire, New York, Ohio, Pennsylvania, Vermont, Virginia and Wisconsin). Regional influenza activity was reported by Guam and 19 states (Alaska, Arizona, Arkansas, California, Idaho, Iowa, Kansas, Michigan, Minnesota, Missouri, Montana, Nevada, New Jersey, New Mexico, North Dakota, Oklahoma, South Carolina, South Dakota and Utah). Local influenza activity was reported by the District of Columbia and 13 states (Alabama, Hawaii, Illinois, Louisiana, Maryland, North Carolina, Oregon, Rhode Island, Tennessee, Texas, Washington, West Virginia and Wyoming). Sporadic influenza activity was reported by the U.S. Virgin Islands and four states (Florida, Georgia, Indiana and Mississippi). Both national and state data are provisional and subject to change as additional reports are received.

Delaware Influenza Surveillance 2015-2016:

During Week 15 there were 101 laboratory-confirmed cases of influenza reported among Delaware residents. Reports of influenza-like illness (ILI) received from participating providers, facilities and institutions in **Delaware show ILI is 0.33%**, (baseline is 1.8%). **Nationally, ILI is 2.1%** (national baseline is 2.1%).

Level of Influenza Activity in Delaware, MMWR Week 15:

Widespread Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in at least half the regions of the state with recent laboratory evidence of influenza in the state.

Influenza-like illness (ILI) is defined as patients presenting with fever of 100° F or greater, cough and/or sore throat in the absence of a known cause other than influenza.

No Activity: No laboratory-confirmed cases of influenza and no reported increase in the number of cases of ILI.

Sporadic: Small numbers of laboratory-confirmed influenza cases or a single laboratory-confirmed influenza outbreak has been reported, but there is no increase in cases of ILI.

Local: Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in a single region of the state.

Regional: Outbreaks of influenza or increases in ILI and recent laboratory-confirmed influenza in at least two but less than half the regions of the state with recent laboratory evidence of influenza in those regions.³

Widespread: Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in at least half the regions of the state with recent laboratory evidence of influenza in the state.

³ Region = population under surveillance in a defined geographical subdivision of a state. Regions typically include several counties. Regional doesn't apply to states with < four counties.

In this report: page Influenza positive cases reported statewide and county, by subtype, Delaware 2015-16 2 2 Influenza positive cases reported statewide and county by age group, Delaware 2015-16 Confirmed cases of influenza by type and subtype, Delaware 2015-16 3 Influenza-related hospitalizations, statewide and county, by age group, Delaware 2015-16 3 Influenza-related deaths, Delaware 2015-16 3 Annual number of influenza cases reported by flu season, Delaware 2004-2005 through 2015-16 4 Percentage of visits for influenza-like illness reported by sentinel providers participating in the ILINet, Delaware 2015-16 4 5 Influenza-like illness reported by ILI reporting partners, Delaware 2015-16 Percentage of Delaware emergency department visits related to influenza, Delaware 2015-16 5

Summary of International Influenza Activity

6

¹ 2015-2016 Region 3 (DE, DC, MD, PA, VA and WV) baseline = 1.8%.

² Laboratory-confirmed case = case confirmed by viral culture or PCR.

								001	awa											
	onfirmed Flu s by Subtype / Lineage	Wk 40- 52	Wk 1	Wk 2	Wk 3	Wk 4	Wk 5	Wk 6	Wk 7	Wk 8	Wk 9	Wk 10	Wk 11	Wk 12	Wk 13	Wk 14	Wk 15	YTD	YTD Total	YTD County %
	A / 2009 H1N1	1	1	0	2	2	2	9	24	58	77	91	67	65	27	15	10	451		
	A / 2012 H3N2	6	0	2	0	0	0	0	1	2	3	2	0	2	1	0	0	19		
Ы	A / no subtype	3	3	0	0	2	2	7	10	38	91	262	169	136	97	50	36	906		
STATEWIDE	A,B co-infection	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	2	1,841	
TAT	B / Yamagata	3	0	0	0	1	0	1	3	3	2	5	6	3	5	3	7	42		
ò	B / Victoria	0	0	0	1	0	2	2	1	4	7	5	9	7	5	4	4	51		
	B / no lineage	4	0	1	0	1	0	1	3	13	17	51	57	60	58	60	44	370		
	A / 2009 H1N1	1	1	0	0	1	0	8	14	42	61	73	52	51	17	14	6	341		
	A / 2012 H3N2	5	0	1	0	0	0	0	1	1	3	0	0	1	1	0	0	13	1,212	65.8%
∕ tle	A / no subtype	1	2	0	0	1	2	2	3	21	64	171	127	86	70	33	18	601		
New Castle County	A,B co-infection	0	0	0	0	0	0	0	0	0	1*	0	0	0	0	0	0	1		
န္ ပိ	B / Yamagata	2	0	0	0	1	0	0	3	3	2	4	6	1	2	2	5	31		
z	B / Victoria	0	0	0	0	0	2	2	0	2	1	3	3	3	0	1	0	17		
	B / no lineage	3	0	0	0	0	0	0	1	7	7	25	30	36	34	39	26	208		
	A / 2009 H1N1	0	0	0	1	1	0	0	5	5	8	5	6	5	3	0	1	40		21.0%
	A / 2012 H3N2	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	387	
Kent County	A / no subtype	2	0	0	0	1	0	3	5	8	19	60	28	32	19	11	14	202		
ΣÖ	B / Yamagata	0	0	0	0	0	0	0	0	0	0	1	0	1	2	1	1	6		
	B / Victoria	0	0	0	1	0	0	0	0	2	0	0	3	1	0	2	2	11		
	B / no lineage	1	0	1	0	0	0	1	2	4	5	20	22	20	17	19	15	127	ļ	
	A / 2009 H1N1	0	0	0	1	0	2	1	5	11	8	13	9	9	7	1	3	70		
	A / 2012 H3N2	1	0	1	0	0	0	0	0	1	0	2	0	0	0	0	0	5		
ty ex	A / no subtype	0	1	0	0	0	0	2	2	9	8	31	14	18	8	6	4	103	242	13.2%
Sussex County	A,B co-infection	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1		
00	B / Yamagata	1	0	0	0	0	0	1	0	0	0	0	0	1	1	0	1	5		
	B / Victoria	0	0	0	0	0	0	0	1	0	6	2	3	3	5	1	2	23		
	B / no lineage	0	0	0	0	1	0	0	0	2	5	6	5	4	7	2	3	35		

Table 1a. Influenza positive¹ cases reported² statewide and county by subtype (A) or lineage (B)³, Delaware 2015-16

- being analyzed at CDC ¥


Table 1b. Influenza positive¹ cases reported² statewide and county by age group, Delaware 2015-16

Cas	nfirmed Flu ses by Age Group	Wk 40- 52	Wk 1	Wk 2	Wk 3	Wk 4	Wk 5	Wk 6	Wk 7	Wk 8	Wk 9	Wk 10	Wk 11	Wk 12	Wk 13	Wk 14	Wk 15	YTD	YTD Total	YTD County %
	0-4 years	2	0	0	2	1	0	5	7	22	41	76	55	69	36	31	20	367		
STATEWIDE	5-24 years	6	2	1	0	1	3	7	8	34	56	131	105	93	47	37	40	571		
1 E	25-49 years	6	1	1	1	1	2	5	15	37	63	103	79	53	58	30	23	478	1,841	
TA.	50-64 years	1	1	0	0	2	1	1	9	15	24	65	39	35	26	24	13	256		
S	65+ years	2	0	1	0	1	0	2	3	10	14	41	31	23	26	10	5	169		
	0-4 years	2	0	0	0	0	0	3	5	13	35	65	45	51	28	27	13	287	1,212	
< tle	5-24 years	4	2	0	0	1	3	4	4	23	35	91	68	61	27	22	23	368		65.8%
ew Cast County	25-49 years	4	0	0	0	1	1	3	5	23	41	51	48	26	33	17	8	261		
New Castle County	50-64 years	0	1	0	0	1	0	1	5	9	17	43	32	24	14	16	7	170		
z	65+ years	2	0	1	0	0	0	1	3	8	11	26	25	16	22	7	4	126		
	0-4 years	0	0	0	2	1	0	1	1	3	5	4	6	7	4	4	4	42	387	21.0%
	5-24 years	1	0	1	0	0	0	1	2	5	7	26	25	23	11	14	13	129		
Kent County	25-49 years	2	0	0	0	0	0	2	5	8	13	30	18	21	14	9	12	134		
δ¥	50-64 years	0	0	0	0	0	0	0	4	2	5	17	6	4	9	4	3	54		
	65+ years	0	0	0	0	1	0	0	0	1	2	9	4	5	3	2	1	28		
	0-4 years	0	0	0	0	0	0	1	1	6	1	7	4	11	4	0	3	38	242	
ty ex	5-24 years	1	0	0	0	0	0	2	2	6	14	14	12	9	9	1	4	74		
Sussex County	25-49 years	0	1	1	1	0	1	0	5	6	9	22	13	6	11	4	3	83		13.2%
<i>s</i> 0	50-64 years	1	0	0	0	1	1	0	0	4	2	5	1	7	3	4	3	32		
	65+ years	0	0	0	0	0	0	1	0	1	1	6	2	2	1	1	0	15		

 ¹ Based on patients with positive nucleic acid or viral culture test results reported to the Division of Public Health.
² Reports are by the date the laboratory results are obtained. As a result, prior weeks' counts may be adjusted to reflect additional cases received. ³ The Division of Public Health Laboratory now has the capability to identify lineage for Influenza B. Since some laboratories

in the state do not have this capability, those influenza cases will be categorized as Influenza B, no lineage identified.

MMWR Week 15 = April 10 - 16, 2016



1 - Based on patients with positive nucleic acid or viral culture test results reported to the Delaware Division of Public Health.

											-	-								
Cas	bitalized Flu ses by Age Group	Wk 40- 52	Wk 1	Wk 2	Wk 3	Wk 4	Wk 5	Wk 6	Wk 7	Wk 8	Wk 9	Wk 10	Wk 11	Wk 12	Wk 13	Wk 14	Wk 15	YTD	YTD Total	YTD County %
	0-4 years	0	0	0	0	0	0	0	1	2	6	6	9	5	2	3	5	39		
STATEWIDE	5-24 years	1	0	0	0	0	0	1	1	1	7	12	5	5	2	3	6	44		
Ē	25-49 years	0	0	0	0	0	0	2	1	3	9	10	5	7	8	5	2	52	317	
STA	50-64 years	0	0	0	0	0	0	1	3	6	11	22	14	14	7	8	0	86		
.,	65+ years	2	0	1	0	1	0	1	1	7`	11	19	20	11	15	5	2	96		
	0-4 years	0	0	0	0	0	0	0	1	0	6	5	9	5	1	3	3	33	254	
v itle	5-24 years	1	0	0	0	0	0	1	1	1	4	10	4	5	2	3	5	37		80.1%
ew Castle County	25-49 years	0	0	0	0	0	0	1	1	2	7	9	5	6	5	4	2	42		
New	50-64 years	0	0	0	0	0	0	1	2	3	7	19	11	11	5	5	0	64		
z	65+ years	2	0	1	0	0	0	1	1	6	8	14	18	8	14	4	1	78		
	0-4 years	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	2	4	41	12.9%
	5-24 years	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	1	3		
Kent County	25-49 years	0	0	0	0	0	0	1	0	1	2	0	0	1	2	1	0	8		
S Y	50-64 years	0	0	0	0	0	0	0	1	1	2	2	3	1	2	2	0	14		
	65+ years	0	0	0	0	1	0	0	0	0	2	4	2	2	0	0	1	12		
	0-4 years	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	2		
nty se	5-24 years	0	0	0	0	0	0	0	0	0	2	1	1	0	0	0	0	4	22	6.9%
Sussex County	25-49 years	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	2		
ω O	50-64 years	0	0	0	0	0	0	0	0	2	2	1	0	2	0	1	0	8		
	65+ years	0	0	0	0	0	0	0	0	1	1	1	0	1	1	1	0	6		

Table 2. Influenza-related hospitalizations statewide and county, by age group,Delaware 2015-16

Table 3. Influenza-related deaths, Delaware 2015-16


ĺ	Influenza- related	Wk 40- 52	Wk 1	Wk 2	Wk 3	Wk 4	Wk 5	Wk 6	Wk 7	Wk 8	Wk 9	Wk 10	Wk 11	Wk 12	Wk 13	Wk 14	Wk 15	YTD
	Deaths	0	0	0	0	0	0	0	0	0	0	0	1	2	0	1	0	4


Influenza Season	Total Annual Influenza Cases										
2004 - 2005	995										
2005 – 2006	541										
2006 - 2007	508										
2007 – 2008	1,401										
2008 – 2009	738										
2009 - 2010	2,247										
2010 – 2011	1,479										
2011 – 2012	267										
2012 – 2013	1,781										
2013 – 2014	1,842										
2014 – 2015	2,390										
2015 – 2016 (YTD)	1,841										

Table 4. Annual number of influenza cases reported by flu season, Delaware 2004-05 through 2015-16

U.S. Outpatient Influenza-Like Illness Surveillance Network (ILINet) Sentinel Providers

An ILINet (sentinel) provider conducts surveillance for influenza-like illness (ILI) in collaboration with the Division of Public Health and the Centers for Disease Control and Prevention (CDC). Data reported by ILINet providers, in combination with other influenza surveillance data, provide a national and statewide picture of influenza activity in the U.S.






¹ 11 of 17 sentinel providers reported.

² Regional baseline is calculated by CDC using non-influenza weeks from the previous three influenza seasons.
Delaware is in Region 3 that includes DE, DC, MD, PA, VA and WV.

Figure 3. Percentage of influenza-like illness (ILI) reported by influenza surveillance ILI reporting partners¹, Delaware 2015-16



¹ ILINet reporting partners include long-term care facilities, colleges / universities, correctional facilities and daycare facilities. Federally gualified health centers are now CDC ILINet sentinel providers.

% ILI= percentage of residents with ILI symptoms. Six long-term care facilities reported.

% ILI= percentage of residents with LL symptome. Survey at 12
3 % ILI= percentage of student visits for ILI; Three universities reported.
4 % ILI= percentage of visits for ILI at the correctional facility; Ten correctional facilities reported.

 $\frac{1}{5}$ % ILI= percentage of children absent with ILI; No daycare providers reported.




Figure 4. Percentage of emergency department visits due to ILI/Flu by MMWR week, Delaware, 2015-16

Summary of International Influenza Activity

In the Northern Hemisphere influenza activity was decreasing, while still elevated in some areas, due in part to an increase of influenza B activity. In the Southern Hemisphere influenza activity was reported to be slightly increasing.

In North America, decreasing influenza activity continued, however high levels remained. Co-circulation of influenza A (predominantly influenza A(H1N1)pdm09) and influenza B was observed. The majority of indicators related to influenza surveillance began to show a decreasing trend in the last several weeks in Canada, Mexico and the United States.

In Central America and the Caribbean, low influenza activity was reported in most countries except in Jamaica where elevated severe acute respiratory infection (SARI) activity associated with influenza A(H1N1)pdm09 virus infection was reported.

In Europe, a decreasing trend of influenza activity was observed overall. In Northern Europe, overall influenza activity decreased but remained at moderate levels. A shift towards circulation of influenza B virus was detected in parts of Europe.

Northern Temperate Asia continued to report ongoing and elevated levels of influenza activity with increasing proportions of influenza B virus.

In tropical South America, low but increasing circulation of influenza A(H1N1)pdm09 virus was reported. In Brazil, influenza activity was above expected levels for this time of year with influenza A(H1N1)pdm09 virus predominating. Colombia reported high circulation of respiratory syncytial virus (RSV). In temperate South America, influenza activity slightly increased but remained at low level. An increase in influenza-like illness (ILI) and SARI rates were reported in Argentina, Chile and Paraguay.

In Oceania and South Africa influenza virus activity remained low.

Source: World Health Organization (WHO), 2016. Influenza Update Number 261 (4/18/16 – based on data up to 4/3/16). Retrieved on 4/21/16 from http://www.who.int/influenza/surveillance_monitoring/updates/latest_update_GIP_surveillance/en/ Reports are updated biweekly.

NOTE: The data provided do not reflect the total number of individuals who have been infected with the influenza virus in Delaware during the reporting period due to the following factors:

- > Many people ill with influenza-like symptoms do not seek medical care.
- > Many who do seek medical care are not tested for influenza.

> The Delaware Public Health Laboratory is limited by capacity to processing a maximum of three specimens per day from each reporting entity.

The Delaware Division of Public Health (DPH) is committed to serving you better by providing the most accurate, upto-date influenza data available.

- For general information on influenza, visit <u>flu.delaware.gov</u> or <u>http://dhss.delaware.gov/dhss/dph/dpc/immunize-</u> flu.html.
- > For specific information on DPH flu clinics, visit <u>http://dhss.delaware.gov/dhss/dph/fluclinics.html</u>.
- For questions on Delaware's weekly flu report, call the DPH Office of Infectious Disease Epidemiology: 302-744-4990.
- > For questions regarding influenza vaccination, please call 302-744-1060.