



Delaware Weekly Influenza Report

MMWR Week 6 (February 2, 2020- February 8, 2020)

Delaware Division of Public Health

National Influenza Synopsis 2019-2020:

National data are updated Friday of each week. Please visit <http://www.cdc.gov/flu/weekly/> for the most current information. Geographic spread of influenza during MMWR Week 1 (February 2, 2020- February 8, 2020) was reported as follows:

- *Sporadic Influenza Activity* was reported in the U.S. Virgin Islands
- *Local Influenza Activity* was reported in the District of Columbia
- *Regional Influenza Activity* was reported in two states (Hawaii and Oregon)
- *Widespread Influenza Activity* was reported Puerto Rico and 48 states (Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming)
- Guam did not report.

Summary of International Influenza Activity:

- In the temperate zone of the northern hemisphere, respiratory illness indicators and influenza activity remained elevated overall.
 - In North America, influenza activity remained elevated influenza A(H1N1)pdm09 and B viruses co-circulating.
 - In Europe, influenza activity continued to increase across the region but appeared to decrease in some countries of Northern Europe.
 - In Central Asia, influenza activity increased with influenza B viruses predominant.
 - In Northern Africa, influenza activity appeared to decrease in Egypt after peaking in recent weeks.
 - In Western Asia, influenza activity remained elevated overall and continued to increase in Lebanon and Turkey.
 - In East Asia, influenza-like illness (ILI) and influenza activity remained elevated overall.
- In the Caribbean and Central American countries, influenza activity was low across reporting countries, except for Mexico with increased activity of influenza A(H1N1)pdm09 viruses. In tropical South American countries, increased influenza activity was reported in Peru.
- In tropical Africa, influenza activity was low across most reporting countries.
- In Southern Asia, influenza activity was low in most reporting countries, but increased in Afghanistan.
- In South East Asia, influenza activity continued to be reported in Lao People's Democratic Republic and Malaysia and increased in Singapore.
- In the temperate zones of the southern hemisphere, influenza activity remained at inter-seasonal levels.
- Worldwide, seasonal influenza A viruses accounted for the majority of detections.

Reference: World Health Organization (WHO), 2017. Influenza Update number 360 (02/03/2020). Retrieved January 20, 2020, from

http://www.who.int/influenza/surveillance_monitoring/updates/latest_update_GIP_surveillance/en/

Reports are updated biweekly.

Delaware Influenza Surveillance 2019-2020:

During MMWR Week 6, there were 785 laboratory-confirmed cases of influenza reported among Delaware Residents. Reports of influenza-like-illness (ILI) received from participating providers, facilities, and institutions in Delaware show ILI is 2.6% compared to Delaware's 2019-2020 baseline of 1.9%. Nationally, 6.8% of visits to a healthcare provider were for ILI, which is an increase from last week, and is still above the 2019-2020 national baseline of 2.4%. All regions remain above their baselines.

Level of Influenza Activity in Delaware, MMWR Week 6:

Widespread:	Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in at least half the regions of the state with recent laboratory evidence of influenza in the state.
Influenza-like illness (ILI) is defined as patients presenting with fever of 100° F or greater, cough and/or sore throat in the absence of a known cause other than influenza.	
CDC Definitions:	
No Activity: No laboratory-confirmed cases of influenza and no reported increase in the number of cases of ILI.	
Sporadic: Small numbers of laboratory-confirmed influenza cases or a single laboratory-confirmed influenza outbreak has been reported, but there is no increase in cases of ILI.	
Local: Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in a single region of the state.	
Regional: Outbreaks of influenza or increases in ILI and recent laboratory-confirmed influenza in at least two but less than half the regions of the state with recent laboratory evidence of influenza in those regions. ³	
Widespread: Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in at least half the regions of the state with recent laboratory evidence of influenza in the state.	

¹ 2018-2019 Region 3 (DE, DC, MD, PA, VA and WV) Baseline = 1.9%.

² Laboratory-confirmed case = case confirmed by viral culture or PCR.

³ Region = population under surveillance in a defined geographical subdivision of a state. Regions typically include several counties. Regional does not apply to states with ≤ four counties.

Summary of Figures for the MMWR Week 6 Report:

Table 1: Influenza Season 2019-2020 – Vaccination Status

Table 1A: Comparison of the 2018-2019 MMWR Week 6 and the 2019-2020 MMWR Week 6 Confirmed Influenza Cases Reported Statewide by County and Age Group

Table 1B: Comparison of the 2018-2019 MMWR Week 6 and the 2019-2020 MMWR Week 6 Confirmed Influenza Cases Reported Statewide by County and subtype/lineage

Figure 1: Confirmed Cases of Influenza by Type and Subtype/Lineage, Delaware 2019-2020 Influenza Season

Table 2: Comparison of the 2018-2019 MMWR Week 6 and the 2019-2020 MMWR Week 6 Influenza-related Hospitalizations and Deaths


Table 3: Annual Number of Influenza Cases Reported by Flu Season, Delaware 2004-05 through 2019-20

Figure 2: Percentage of Visits for Influenza-Line Illness Reported by Sentinel Providers participating in the U.S. Outpatient ILI Surveillance Network (ILINet), Delaware 2019-2020

Figure 3A: Percentage of Emergency Department (ED) Visits Due to ILI/Influenza by MMWR Week, Delaware 2019-2020

Figure 3B: County-Specific Percentage of ED visits Due to ILI/Influenza by MMRW Week, Delaware 2019-2020

Table 1: Influenza Season 2019-2020 – “Vaccination Status of Confirmed Cases”



*Received influenza vaccination this season per DelVax
†Not vaccinated for influenza this season per DelVax
¶No record in the Delaware Immunization Registry (DelVax)

Table 1A: Comparison of the 2018-2019 MMWR Week 6 and the 2019-2020 MMWR Week 6 Confirmed¹ Influenza Cases Reported Statewide by County and Age Group

Confirmed Flu Cases by Age Group		2018-2019 Influenza Season					2019-2020 Influenza Season				
		Week 5	Total Counts	County Percentage (%)	YTD ²	YTD County Percentage (%)	Week 5	Total Counts	County Percentage (%)	YTD ²	YTD County Percentage (%)
STATEWIDE	0-4	121	630	--	2694	--	172	785	--	4383	--
	5-9	113					134				
	10-14	90					76				
	15-19	23					34				
	20-29	5					71				
	30-39	50					84				
	40-59	85					118				
	60+	98					96				
New Castle County	0-4	58	259	41.11	1104	40.98	92	354	45.1	1994	45.49
	5-9	38					47				
	10-14	32					90				
	15-19	9					9				
	20-29	18					2				
	30-39	22					42				
	40-59	38					61				
	60+	44					41				
Kent County	0-4	34	166	26.35	680	25.24	30	168	21.53	1078	24.60
	5-9	29					30				
	10-14	8					15				
	15-19	10					6				
	20-29	18					25				
	30-39	16					13				
	40-59	20					28				
	60+	21					22				
Sussex County	0-4	29	205	32.54	910	33.78	49	249	31.272	1298	29.61
	5-9	46					53				
	10-14	40					30				
	15-19	4					19				
	20-29	14					13				
	30-39	12					26				
	40-59	27					27				
	60+	33					32				

¹Influenza Cases are confirmed via PCR testing

²YTD stands for “Year to Date” and represents the cumulative number of cases through the current MMWR Week being assessed for the 2018-19 and 2019-20 influenza seasons, respectively.

Table 1B: Comparison of the 2018-2019 MMWR Week 6 and the 2019-2020 MMWR Week 6 Confirmed¹ Influenza Cases Reported Statewide by County and subtype/lineage

Confirmed Flu Cases by Subtype / Lineage		2018-2019 Influenza Season		2019-2020 Influenza Season	
		Week 6	Total Counts	Week 6	Total Counts
STATEWIDE	A / 2009 H1N1	38	630	56	785
	A / 2012 H3N2	34		0	
	A / no subtype	539		412	
	Co-infection	0		1	
	B / Yamagata	0		0	
	B / Victoria	0		14	
	B / no lineage	17		302	
New Castle County	A / 2009 H1N1	24	259	40	354
	A / 2012 H3N2	26		0	
	A / no subtype	209		207	
	Co-infection	0		0	
	B / Yamagata	0		0	
	B / Victoria	0		11	
	B / no lineage	0		96	
Kent County	A / 2009 H1N1	14	166	11	169
	A / 2012 H3N2	6		0	
	A / no subtype	133		80	
	Co-infection ²	0		1	
	B / Yamagata	0		0	
	B / Victoria	0		0	
	B / no lineage	13		77	
Sussex County	A / 2009 H1N1	0	205	5	249
	A / 2012 H3N2	4		0	
	A / no subtype	197		122	
	Co-infection	0		0	
	B / Yamagata	0		0	
	B / Victoria	0		3	
	B / no lineage	4		119	

¹Influenza Cases are confirmed via PCR testing

Confirmed Cases of Influenza by Type and Subtype/Lineage, Delaware 2019-2020 Influenza Season




Figure 1: During MMWR Week 6 for the 2019-2020 Influenza season, there were 32 confirmed cases of Influenza reported by ILINet providers. Nine were identified as subtype B/Victoria, and Twenty-three were identified as Flu A/ H1N1. The percent positive for samples tested was 52.4 %.

¹Percent Positive represents the amount of specimens that tested positive for influenza out of all samples tested for that MMWR week.

Table 2: Comparison of the 2018-2019 MMWR Week 6 and the 2019-2020 MMWR Week 6 Influenza-related Hospitalizations and Deaths Statewide

Hospitalizations and Deaths due to Influenza	2018-2019 Influenza Season				2019-2020 Influenza Season			
	Week 6	YTD Totals ¹	Percentage of Confirmed Cases (%) ²	YTD Percentage of Confirmed Cases (%) ³	Week 6	YTD Totals ¹	Percentage of Confirmed Case (%) ²	YTD Percentage of Confirmed Cases (%) ³
Hospitalizations	86	441	13.6	16.6	43	240	5.85	5.47
Deaths	0	6	0	.22	0	7	0	0.15

¹YTD stands for “Year to Date” and represents the cumulative number of cases through the current MMWR Week that were hospitalized or died

²Percentage of cases confirmed during the single MMWR Week

³Percentage of cases for the cumulative count of confirmed cases through the influenza season to the current MMWR Week.

Table 3: Annual Number of Influenza Cases Reported by Flu Season, Delaware 2004-05 through 2019-20

Influenza Season	Total Annual Influenza Cases
2004 – 2005	995
2005 – 2006	541
2006 – 2007	508
2007 – 2008	1,401
2008 – 2009	738
2009 – 2010	2,247
2010 – 2011	1,479
2011 – 2012	267
2012 – 2013	1,781
2013 – 2014	1,842
2014 – 2015	2,390
2015 – 2016	1,842
2016 – 2017	4,590
2017 – 2018	9,050
2018 – 2019	6,387
2019 – 2020 (YTD)	4383

U.S. Outpatient Influenza-Like Illness Surveillance Network (ILINet) Sentinel Providers

An ILINet (sentinel) provider conducts surveillance for influenza-like illness (ILI) in collaboration with the Division of Public Health and the Centers for Disease Control and Prevention (CDC). Data reported by ILINet providers, in combination with other influenza surveillance data, provide a national and statewide picture of influenza activity in the U.S.

Percentage of Visits for Influenza-Like Illness Reported by Sentinel Providers¹ participating in the U.S. Outpatient ILI Surveillance Network (ILINet), Delaware 2019-2020




Figure 2: Delaware's regional baseline² for healthcare visits relating to ILI symptoms in the 2019-2020 Influenza Season is 1.9% and the national baseline³ is 2.4%. In MMWR Week 6, the amount of ILI related visits reported by sentinel providers was inconclusive for Week 6.

¹seven of twelve sentinel providers reported

²The regional baseline is calculated by the CDC using non-influenza weeks from the previous three influenza seasons. Delaware is in Region 3, which also includes DC, MD, PA, VA, and WV.

³The National baseline is calculated by the CDC using non-influenza weeks from the previous three influenza seasons.

Percentage of Emergency Care Visits Due to ILI/Influenza by MMWR Week, Delaware 2019-2020




Figure 3A: Syndromic data collected from ESSENCE shows that in the weeks preceding the start of the 2019-2020 Influenza season and through MMWR Week 6, there is an increasing trend in the percentage of ED visits due to ILI/Flu symptoms.

County-Specific Percentage of ED visits Due to ILI/Influenza by MMRW Week, Delaware
2019-20




Figure 3B: Syndromic data collected from ESSENCE shows that in the weeks preceding the start of the 2019-2020 Influenza season and through MMWR Week 6 stratified by Delaware’s three counties.

NOTE: Data provided do not reflect the total number of individuals who have been infected with the influenza virus in Delaware during the reporting period due to the following factors:

- Many people ill with influenza-like symptoms do not seek medical care.
- Many who do seek medical care are not tested for influenza.
- The Delaware Public Health Laboratory is limited by capacity to processing a maximum of three specimens per day from each reporting entity.

The Delaware Division of Public Health (DPH) is committed to serving you better by providing the most accurate, up-to-date influenza data available.

- For general information on influenza, visit flu.delaware.gov or <http://dhss.delaware.gov/dhss/dph/dpc/immunize-flu.html>.
- For specific information on DPH flu clinics, visit <http://dhss.delaware.gov/dhss/dph/fluclinics.html>.
- For questions on Delaware’s weekly flu report, call the DPH Office of Infectious Disease Epidemiology at 302-744-4990.
- For questions regarding influenza vaccination, please call 302-744-1060.