



ANESTHESIA INFECTION PREVENTION CHECKLIST

OR: _____ SUITE _____ OBSERVER: _____
 PROVIDER NAME: _____ DATE: _____ TIME: _____

	Self Assessment		Pass	Fail	Re-Check
	Pass	Fail			
1. Hand washing performed before first case. Surgical scrub performed before first case using Chlorhexidine Gluconate (CHG) is preferred but not mandatory.					
2. Hand hygiene performed prior to accessing/preparing clean supplies/medications: i.e. preparing syringes, drawing up medications, accessing medication vials.					
3. Hand hygiene performed prior to gloving and after removing gloves.					
4. Hand hygiene performed between dirty and clean steps of a procedure: i.e.:suctioning a patient followed by administering IV medications.					
5. Full barrier protection is worn for central line insertions and Central Line bundle is followed and documented (see attached copy of Bundle sheet).					
6. Gloves are worn when inserting IVs (see #3 above).					
7. Laryngeal Mask Airway (LMA) / laryngoscopes are handled with gloves (see #3 above).					
8. Gloves are removed after inserting or removing LMA / laryngoscope.					

Soap and water must be used for hand hygiene if hands are bloody or soiled; otherwise, alcohol hand rubs are preferred. CHG containing hand scrub or regular hand scrub are acceptable.

We appreciate your input.

Please send comments to: Hpxton@che-east.org.
 Thank you.